

PREPARED FOR: AURORA MOLD INSPECTIONS

TEST ADDRESS: 123 MAIN STREET ANYTOWN, NJ 07676

CERTIFICATE OF MOLD ANALYSIS

PREPARED FOR:

AURORA MOLD INSPECTIONS

PHONE NUMBER: (201) 476-9200

EMAIL: LARRY@AURORAHOMEINSPECTIONS.COM

TEST LOCATION:

JANE AND JOE HOMEOWNER

123 MAIN STREET

ANYTOWN, NJ 07676

CHAIN OF CUSTODY # 52034724

COLLECTED: MON APRIL 13, 2020

RECEIVED: MON APRIL 13, 2020

REPORTED: MON APRIL 13, 2020



APPROVED BY: JOHN D. SHANE PH.D.,
LABORATORY MANAGER

VERSION: 1.0 (A VERSION NUMBER GREATER THAN ONE (1) INDICATES THAT THE DATA IN THIS REPORT HAS BEEN AMENDED)

EPA regulations or standards for airborne or surface mold concentrations have not been established. There are also no EPA regulations or standards for evaluating health effects due to mold exposure. Information about mold can be found at www.epa.gov/mold.

All samples were received in an acceptable condition for analysis unless noted specifically in the Comments section under a particular sample. All results relate only to the samples submitted for analysis and apply to the samples as received by the laboratory. Volumes, flowrates, areas or other information are supplied by the customer. This information can affect the validity of the results. Results have not been adjusted for field or laboratory unless otherwise noted. InspectorLab bears no responsibility for sample collection activities or analytical method limitations. No warranty is either express or implied and InspectorLab assumes no responsibility or liability for error in public information utilized, statements from sources other than InspectorLab, or developments resulting from situations outside the scope of this analysis, nor for the purpose for which the client uses the analysis. The determinations in this report are outside the scope of the AIHA LAP, LLC scope of accreditation. Contractors or consultants reviewing this report must draw their own conclusions regarding further investigation or remediation deemed necessary. InspectorLab liability is limited to the cost of the sample analysis and may not exceed the amount of the fee paid by the client.

Reports are issued free of alterations or additions and InspectorLab does not accept liability of the tampering or unlawful alteration of documents sent. All reports are expressly and exclusively for InspectorLab clients and may not be reproduced by third parties. If this report is reproduced, it must be reproduced in full unless written permission is obtained from InspectorLab. InspectorLab keeps all client data secure and confidential and any information contained in reports or files will not be divulged unless permission is expressly given by the client submitting the sample(s) except where authorized by law and all InspectorLab employees are required to maintain the confidentiality of all non-public personal information provided. We do not sell client information to anyone or disclose client information to marketing companies. This disclaimer governs the use of this report. By using or accepting this report, you accept this disclaimer in full.

Detailed Mold Report

(WATER-INDICATING FUNGI, IF PRESENT, ARE SHOWN BELOW IN RED)

Analysis Method	Air Analysis	Air Analysis	Air Analysis	Surface Analysis
Lab Sample #	52034724-1	52034724-2	52034724-3	52034724-4
Sample Identification	21023444	21045322	21042361	21042362
Sample Location	OUTSIDE CONTROL	MASTER BEDROOM	TV ROOM	TV ROOM CLOSET WALL
Sample Type / Metric	Air-O-Cell/150L	Air-O-Cell/150L	Air-O-Cell/150L	Swab
Analysis Date	Wed June 10, 2015	Wed June 10, 2015	Wed June 10, 2015	Wed June 10, 2015
Determination	CONTROL	NORMAL	PROBLEM	GROWTH

Fungal Types Identified	Raw Count	Spores / m ³	% of Total	Raw Count	Spores / m ³	% of Total	Raw Count	Spores / m ³	% of Total	Mold Present	
*INDOOR PROBLEM FUNGI											
Cladosporium sphaerospermum	---	---	---	---	---	---	---	---	---	X	
Penicillium	---	---	---	---	---	---	---	---	---	X	
Penicillium/Aspergillus	---	---	---	---	---	---	1,625	10,888	66	X	
Scopulariopsis	---	---	---	---	---	---	771	5,166	31	X	
**Non-Problem Fungi											
Alternaria	7	47	2	---	---	---	3	20	<1	---	
Ascospores	19	127	7	17	114	9	---	---	---	---	
Basidiospores	47	315	17	7	47	4	9	60	<1	---	
Bipolaris/Drechslera	5	34	1	1	7	<1	---	---	---	---	
Cladosporium	124	831	47	37	248	21	3	20	<1	---	
Curvularia	7	47	2	1	7	<1	---	---	---	---	
Epicoccum	6	40	2	2	13	1	---	---	---	---	
Nigrospora	4	27	1	---	---	---	---	---	---	---	
Penicillium/Aspergillus	6	40	2	91	610	52	*	*	*	*	
Pithomyces	6	40	2	2	13	1	---	---	---	---	
Smut/Myxomycetes	31	208	11	14	94	8	17	114	<1	---	
Total Spore Count	260	1,800	100	170	1,200	100	2,400	16,000	100	NA	
Minimum Detection Limit	7			7			7			1	
Comments/Definitions	<p>CONTROL samples are normally taken outside a building to provide a baseline from which samples on the interior of the building are compared. Outside air is considered normal whatever the mold counts may be. Light Debris: The debris present in the sample likely had no effect on the accuracy of the mold count.</p>			<p>Mold counts are within a NORMAL RANGE and there is no indication, based on the mold counts, that there is any exposure concern to the occupants. The LIGHT DEBRIS present in the sample likely had no effect on the accuracy of the mold count.</p>			<p>Mold concentrations in the air are ABNORMAL and based on the mold counts, you likely have a mold source from which spores are able to become airborne and are an exposure concern to the occupants. MODERATE DEBRIS: in the sample likely had limited affect on the accuracy of the mold count.</p>			<p>Presence of current or former MOLD GROWTH observed. EXPOSURE TO SPORES LIKELY and will continue if the growth is not removed. An active or intermittent water source will cause the mold to continue to grow if the water source is not eliminated.</p>	

* Indoor Problem Fungi are generally capable of growing on wetted building materials.

** Non-Problem Fungi are less capable or do not grow on wetted building materials. They are commonly found in the air outside and infiltrate into indoor air naturally. High numbers of any one of these spore types as compared to the Control sample may indicate that they are growing on wetted building materials indoors.

Spore types not listed in this report were not observed.

Background debris estimates the amount of non-spore particles. Increasing amount of debris will affect the accuracy of the spore counts. Total percent may not equal 100% due to rounding.

Total Spore Counts are reported to 2 significant figures.

PREPARED FOR: AURORA MOLD INSPECTIONS

TEST ADDRESS: 123 MAIN STREET ANYTOWN, NJ 07676

Introduction

All spores found in indoor air are also normally found in outdoor air because most originate or live in the soil and on dead or decaying plants. Therefore, it is not unusual to find mold spores in indoor air. This Mold Glossary is only intended to provide general information about the mold found in the samples that were provided to the laboratory.

Alternaria

Outdoor Habitat: One of the most commonly observed spores in the outdoor air worldwide, normally in low numbers.

Indoor Habitat: Capable of growing on a wide variety of substrates and manufactured products found indoors when wetted.

Allergy Potential: Type I (hay fever, asthma), Type III (hypersensitivity pneumonitis), Common cause of extrinsic asthma

Disease Potential: Not normally considered a pathogen, but can become so in immunocompromised persons.

Toxin Potential: Several known

Comments: One of the most common and potent allergens in the indoor and outdoor air. Seen in indoor air in low concentrations, probably as a result of outdoor air infiltration and/or recycling of settled dust.

Ascospores

Outdoor Habitat: Soil and decaying vegetation, dead and dying insects. These spores constitute a large part of the spores in the air and can be found in the air in very large numbers in the spring and summer, especially during and up to three (3) days after a rain.

Indoor Habitat: Very few of fungi that produce ascospores grow indoors. Some fungi that produce ascospores are recognizable by their spores and when observed are listed under their own categories. Wetted wood and gypsum wallboard paper

Allergy Potential: Depends on the type of fungus producing the ascospores.

Disease Potential: Not normally pathogenic as a group

Toxin Potential: None known

Comments: Ascospores are produced from a very large group of fungi. Notable ascospores that are considered problematic for indoor environments are Chaetomium, Peziza, and Ascotracha. If these types of ascspores are observed they will be listed in the report under their own names.

PREPARED FOR: AURORA MOLD INSPECTIONS

TEST ADDRESS: 123 MAIN STREET ANYTOWN, NJ 07676

Basidiospores

Outdoor Habitat: These are mushroom spores and are common everywhere outside, especially in the late summer and fall.

Indoor Habitat: Mushrooms can grow on very wet wood products, especially on footer plates, basements, and crawlspaces. Sometimes mushrooms can be observed growing in potted plants indoors.

Allergy Potential: Rarely reported, but some Type I (hay fever, asthma) and Type III (hypersensitivity pneumonitis) has been reported.

Disease Potential: None known

Toxin Potential: None known

Comments: Mushroom spores are commonly found indoors, especially when the outdoor spore count is high. When spores of this group are derived from wood rotting fungi, including dry rot (*Serpula* and *Poria*), they can be especially destructive to buildings. When spores from destructive types of mushrooms (dry and wet rot group) are observed in the sample they are listed under their own names on the report.

Bipolaris/Drechslera

Outdoor Habitat: Commonly observed spores in the outdoor air worldwide, normally in low numbers.

Indoor Habitat: Wetted wood and gypsum wallboard paper

Allergy Potential: Type I (hay fever, asthma)

Disease Potential: Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals.

Toxin Potential: None known

Comments: This category represents at least three genera, including *Bipolaris*, *Drechslera*, and *Exserohilum*. This group cannot be consistently separated by spore morphology alone.

PREPARED FOR: AURORA MOLD INSPECTIONS

TEST ADDRESS: 123 MAIN STREET ANYTOWN, NJ 07676

Cladosporium

Outdoor Habitat: Cladosporium is one of the most common environmental fungi observed worldwide and is widely reported from soil and decaying vegetation.

Cladosporium herbarum and C. cladosporioides are among the most frequently encountered species, both in outdoor and indoor environments.

Indoor Habitat: Wetted wood and gypsum wallboard paper, paper products, textiles, rubber, window sills. Cladosporium has the ability to grow at low temperatures and can thus, grow on rubber gaskets and food in refrigerators.

Allergy Potential: Type I (hay fever, asthma) - an important and common outdoor allergen

Disease Potential: Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals. Cladosporium are some of the most common species reported as indoor contaminants, occasionally linked to health problems.

Toxin Potential: Cladosporium has two known toxins (cladosporin and emodin). These toxins are not known to be highly toxic. There is no evidence in the literature of toxic effects associated to inhalation of Cladosporium conidia (spores) indoors.

Comments: The most commonly reported spore in the outdoor air worldwide. This makes Cladosporium one of the most commonly reported and abundant spore types both indoors and outdoors. The prevalence of this spore can vary throughout the year, but is especially high in late summer and autumn, especially where cereal crops are commonly planted.

An important and common allergen source.

Cladosporium sphaerospermum

Outdoor Habitat: Dead or dying cellulosic materials like wood and leaves.

Indoor Habitat: A favorite place for this mold type to grow is on wetted or rotting window sills. Also likes to grow on all type of wetted or rotting wood.

Allergy Potential: Type I (hay fever, asthma) - an important and common outdoor allergen

Disease Potential: Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals.

Toxin Potential: None known

Comments: A common mold on wetted wood, especially on window sills. Not frequently found in the air.

PREPARED FOR: AURORA MOLD INSPECTIONS

TEST ADDRESS: 123 MAIN STREET ANYTOWN, NJ 07676

Curvularia

Outdoor Habitat: Soil and decaying vegetation

Indoor Habitat: Wetted wood and gypsum wallboard paper, many cellulytic substrates

Allergy Potential: Type I (hay fever, asthma), common cause of allergenic rhinitis

Disease Potential: Potential human pathogen in immunocompromised people

Toxin Potential: None known

Comments: None

Epicoccum

Outdoor Habitat: Epicoccum is a widespread cosmopolitan that grows on dead or decaying organic matter, wood, textiles, paper, a variety of foods, insects and human skin. It is commonly found in the soil. Epicoccum spores are more prevalent on dry, windy days, with higher counts late in the day.

Indoor Habitat: Capable of growing on a wide variety of substrates and manufactured products found indoors when wetted such as gypsum board, floors, carpets, mattress dust, and house plants.

Allergy Potential: Type I (hay fever, asthma)

Disease Potential: None known

Toxin Potential: None known

Comments: Very common in outdoor air in the summer months, especially in the midwest USA during harvest times.

Nigrospora

Outdoor Habitat: Soil and decaying vegetation

Indoor Habitat: Wetted wood and gypsum wallboard paper

Allergy Potential: Type I (hay fever, asthma)

Disease Potential: None known

Toxin Potential: None known

Comments: Rarely observed growing indoors

PREPARED FOR: AURORA MOLD INSPECTIONS

TEST ADDRESS: 123 MAIN STREET ANYTOWN, NJ 07676

Penicillium**Outdoor Habitat:** Soil and decaying vegetation, textiles, fruits**Indoor Habitat:** Wetted wood and gypsum wallboard paper, textiles, leather**Allergy Potential:** Type I (hay fever, asthma), Type III (hypersensitivity pneumonitis)**Disease Potential:** Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals.**Toxin Potential:** Several known**Comments:** Extremely common in indoor air, but without the fruiting bodies associated with the spores will be listed as "Penicillium / Aspergillus" group. Penicillium identified in air samples indicates that the fruiting bodies were observed. This usually suggests that the source of the mold is nearby and / or a growth was disturbed. The fruiting bodies are not easily sent airborne, nor do they stay in the air long.***Penicillium/Aspergillus*****Outdoor Habitat:** Soil and decaying vegetation, textiles, fruits. These spores are commonly observed and are a normal part of outside air.**Indoor Habitat:** Wetted wood and gypsum wallboard paper, textiles, leather, able to grow on many types of substrates.**Allergy Potential:** Type I (hay fever, asthma), Type III (hypersensitivity pneumonitis)**Disease Potential:** Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals.**Toxin Potential:** Several known**Comments:** Extremely common in indoor air in low amounts. This type of spore should not constitute an overwhelming percentage and/or be present in very high numbers as compared to the outside (control).

These two genera are grouped together because they cannot be reliably differentiated into their respective genera based solely on spore morphology.

PREPARED FOR: AURORA MOLD INSPECTIONS

TEST ADDRESS: 123 MAIN STREET ANYTOWN, NJ 07676

Pithomyces

Outdoor Habitat: Soil and decaying vegetation and their spores are easily dispersed into the air by wind

Indoor Habitat: Wetted wood and gypsum wallboard paper

Allergy Potential: None known

Disease Potential: None known

Toxin Potential: One known (sporidesmin)

Comments: A very common spore type in the air. Can be a water indicator mold type indoors

Scopulariopsis

Outdoor Habitat: Soil and decaying vegetation, dung

Indoor Habitat: Wetted wood and gypsum wallboard paper

Allergy Potential: Type III (hypersensitivity pneumonitis)

Disease Potential: Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals.

Toxin Potential: Not well studied

Comments: Easily dispersed by wind and air currents. Can grow with very little water and readily grow on wallboard when high humidity situations, e.g. closets - capable of growing on leather clothes

Smut/Myxomycetes

Outdoor Habitat: Soil and decaying vegetation and wood, especially dead stumps and bark

Indoor Habitat: Not known to grow indoors, sometimes found on firewood

Allergy Potential: Type I (hay fever, asthma), rare

Disease Potential: None known

Toxin Potential: None known

Comments: These two groups are difficult to distinguish due to their "round, brown" morphology. Smuts are especially common in the environment and can be seen in indoor air samples even during the winter in homes because the spores can get trapped in carpets
